



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A NEW DERIVATIVE OF DIAZO-AMIDO-BENZENE.

(2, 2' dibrom, 4, 4' dimethyl diazo amido-benzene.)

By EDWARD BARTOW and H. C. ALLEN, University of Kansas, Lawrence.

Read before the Academy, at Topeka, December 31, 1904.

M-BROMTOLUENE may be prepared from *p*-acettoluid by bromination, elimination of the acetyl group, and submitting the amine thus formed to the Griess reaction,¹ thus replacing the amido group with hydrogen.

In following out the Griess reaction for the diazotizing of *m*-brom-*p*-toluidine by means of nitrous acid, we obtained well-defined bright yellow, needle-like crystals. Wroblevsky,² who prepared *m*-bromtoluene by the above-mentioned series of reactions, warms the alcoholic solution after diazotizing to effect reduction and makes no mention of an intermediate product. Sellards³ mentions a crystal broth which is formed before the reduction takes place.

Our method of procedure was as follows: *P*-acettoluid was suspended in water and brominated according to Wroblevsky² by adding bromine with occasional shaking, but without particular care to cool the reaction mixture. Under these circumstances *m*-brom-*p*-acettoluid is formed.

The acetyl group, which serves to protect the amine, is then removed by saponifying with either NaOH or KOH. Wroblevsky² claims that only KOH will act, while Grete⁴ and Sellards³ find that NaOH does equally well. Our experience was that either would react, but that the NaOH required longer heating, and also heating on the sand-bath. The *m*-brom-*p*-toluidine was then distilled with steam, dried with calcium chloride and diazotized by leading nitrous acid into a solution in absolute alcohol. The nitrous acid is best prepared by using arsenic trioxide having a specific gravity of 1.3 to 1.35. The nitrous acid was conducted through a reflux condenser, cooled by ice-water, and a Woulff's flask, surrounded by a freezing mixture, into the alcoholic solution. The condenser and flask serve as traps for nitric acid as well as for cooling the nitrous acid. The gas was absorbed until near the end of the reaction, and at the same time the bright yellow crystals mentioned above were formed.

If any acid is present in the alcoholic solution, the substance formed will be reduced to *m*-bromtoluene. Ten drops of sulfuric acid in ten grams of the substance completely reduced it in thirty minutes. When no acids are present, however, the crystals are very

1. A, 121-258. 2. A, 168-154. 3. Dissertation, U. of K., 1904. 4. A, 177-232.

stable and can be isolated. They are soluble in ether, but only slightly so in alcohol, and are purified by washing with alcohol and recrystallizing from a mixture of ether and alcohol.

A determination for bromine gave the following results :

0.2151 grams of the substance gave 0.2100 grams of AgBr.

Calculated for $C_{14}H_{13}N_3Br_2$.	Found.
41.77%	41.58%



A molecular weight determination, by lowering of the freezing-point of benzene, gave the following :

I. 0.2435 grams caused a depression of 0.223 degrees in the freezing-point of 14.13 grams of benzene.

II. 0.4459 grams caused a depression of 0.42 degrees in the freezing-point of 14.13 grams of benzene.

	Calculated for $C_{14}H_{13}N_3Br_2$.	Found.
I.....	382.13	378.66
II.....	368.16

If, as claimed by Griess, aniline treated in this way gives diazo-amido-benzene, then our reaction may be represented as follows :



The substance may be graphically represented :

